

LOCAL GEOMETRY OF SINGULAR REAL ANALYTIC SURFACES

DANIEL GRIESER

ABSTRACT. Let $V \subset \mathbb{R}^N$ be a compact real analytic surface with isolated singularities, and assume its smooth part V_0 is equipped with a Riemannian metric that is induced from some analytic Riemannian metric on \mathbb{R}^N . We prove:

1. Each point of V has a neighborhood which is quasi-isometric (naturally and 'almost isometrically') to a union of metric cones and horns, glued at their tips.
2. A full asymptotic expansion, for any $p \in V$, of the length of $V \cap \{q : \text{dist}(q, p) = r\}$ as $r \rightarrow 0$.
3. A Gauss-Bonnet Theorem, saying that horns do not contribute an extra term, while cones contribute the leading coefficient in the length expansion of 2.
4. The L^2 Stokes Theorem, self-adjointness and discreteness of the Laplace-Beltrami operator on V_0 , and a Gauss-Bonnet Theorem for the L^2 Euler characteristic.

As a central tool we use resolution of singularities.

1. INTRODUCTION

In this paper we investigate in detail some aspects of the geometry and analysis of a compact real analytic surface with isolated singularities. By this we mean a compact subset V of some \mathbb{R}^N which is locally given as zero set of a finite number of real analytic functions, and such that V is a smooth 2-dimensional manifold outside a finite set $\text{sing } V$. Let $V_0 = V \setminus \text{sing } V$ be the smooth part. In order to speak of geometry or analysis on V , we need to introduce a metric. We assume that an *induced metric* g is given on V , i.e. a Riemannian metric on V_0 which is induced by some real analytic Riemannian metric on \mathbb{R}^N . g induces an (intrinsic) distance function d on V .

Our first concern is a description of the local structure of the metric space (V, d) up to quasi-isometry. Here, two metric spaces (X, d) , (X', d') are called *quasi-isometric* if there is a bijective map $\Phi : X \rightarrow X'$ which is a quasi-isometry, i.e. such that there is a constant $C > 0$ with

$$C^{-1}d(x, y) \leq d(\Phi(x), \Phi(y)) \leq Cd(x, y) \quad \text{for all } x, y \in X.$$

Date: January 1999.

The author gratefully acknowledges support by the Deutsche Forschungsgemeinschaft (Gerhard-Hess-Programm) and the Erwin-Schrödinger Institute.

The model metric spaces that will occur are cones and horns. These are defined as follows: Let $\gamma \geq 1$ and $\varepsilon > 0$. The space

$$C_\gamma = ([0, \varepsilon)_r \times S_\theta^1) / (\{0\} \times S^1)$$

with distance function d_γ induced by the Riemannian metric

$$dr^2 + r^{2\gamma} d\theta^2$$

for $r > 0$ and $d_\gamma(0, (r, \theta)) = r$, where $0 := [\{0\} \times S^1]$ is the 'tip', is called γ -horn and γ its exponent. A 1-horn is also called a cone.

Theorem 1.1. *Let (V, d) be as above. Then each $p \in V$ has a neighborhood (in V) which is quasi-isometric to a union of finitely many cones and horns with rational exponents, glued at their tips.*

The distance function d on the space glued from $C_{\gamma_1}, \dots, C_{\gamma_n}$ is given by the d_{γ_i} on each C_{γ_i} and $d(p, q) = d_{\gamma_i}(p, 0) + d_{\gamma_j}(0, q)$ if $p \in C_{\gamma_i}, q \in C_{\gamma_j}, i \neq j$.

Example If $a \leq b$ are positive integers then the set

$$\{(x, y, z) : (x^2 + y^2)^a = z^{2b}\} \subset \mathbb{R}^3$$

with metric induced from the Euclidean metric on \mathbb{R}^3 is quasi-isometric (near zero) to two b/a -horns glued at their tip. From this example one sees that any combination of cones and horns can occur if one allows V to be *semi*-analytic (i.e. allowing inequalities in a local description of V) or even semi-algebraic, so Theorem 1.1 gives a complete classification in this case.

Note that a small ball around a smooth point is quasi-isometric to a Euclidean ball, which is a cone in polar coordinates.

Theorem 1.1 can be sharpened considerably: First, the quasi-isometry can be chosen to be 'natural', and second, it gets closer and closer to an isometry as one approaches p . Before we state this precisely, we introduce the infinitesimal notion of quasi-isometry: Two Riemannian metrics g, g' on a manifold M are called quasi-isometric if there is $C > 0$ such that for each $x \in M$ one has the inequality of quadratic forms

$$C^{-1}g_x \leq g'_x \leq Cg_x.$$

If this holds, then we write

$$g \approx_C g'$$

or simply $g \approx g'$. C is called the quasi-isometry constant. It is easy to see that, for two connected Riemannian manifolds $(M, g), (N, h)$, a diffeomorphism $\Phi : M \rightarrow N$ is a quasi-isometry for the induced distance functions if and only if $\Phi^*h \approx g$.

Let dist denote the (extrinsic) distance function on \mathbb{R}^N and $B_\varepsilon(p) = \{x \in V : \text{dist}(x, p) < \varepsilon\}$.

Theorem 1.2. *Let (V, g) be as above and let $p \in V$. If ε_0 is sufficiently small then the following holds for each connected component K of $B_{\varepsilon_0}(p) \setminus p$: There is a parametrization*

$$\Phi : (0, \varepsilon_0) \times S^1 \rightarrow K$$

which 'preserves' r , i.e.

$$(1) \quad \text{dist}(\Phi(r, \theta), p) = r \quad \text{for all } r < \varepsilon_0$$

and is normalized arc length along each $K_r := \{q \in K : \text{dist}(p, q) = r\}$ such that

$$(2) \quad \Phi^*g \cong_{1+\delta} dr^2 + \left(\frac{l(r)}{2\pi}\right)^2 d\theta^2.$$

Here, $l(r)$ is the length of K_r . In (2), one can choose

$$\delta = C\varepsilon^\alpha$$

when restricting Φ to $(0, \varepsilon) \times S^1$, $\varepsilon \leq \varepsilon_0$, with C and $\alpha > 0$ only depending on V and g .

Explicitly, the arc length condition on Φ means

$$(3) \quad |\partial\Phi/\partial\theta|_g = l(r)/2\pi.$$

Clearly, a parametrization Φ satisfying (1) and (3) (if it exists) is uniquely determined by one curve $r \mapsto \Phi(r, \theta_0)$ and an orientation on K .

In order to obtain Theorem 1.1 from Theorem 1.2 one needs to know the leading asymptotic behavior of $l(r)$ as $r \rightarrow 0$. The following theorem gives the complete asymptotics.

Theorem 1.3. *The function $l(r)$ defined in Theorem 1.2 has an asymptotic expansion, as $r \rightarrow 0$,*

$$(4) \quad l(r) \sim \sum_{i,j} C_{i,j} r^i (\log r)^j$$

with i ranging over $n/m \geq 1$ for some fixed integer m and arbitrary integers $n \geq m$, and j ranging over $\{0, 1\}$. Also, the leading term contains no logarithm, i.e. if

$$\gamma = \min\{i : C_{i,0} \neq 0 \text{ or } C_{i,1} \neq 0\}$$

then $C_{\gamma,1} = 0$.

The asymptotic expansion (4) may be differentiated (arbitrarily often) term by term.

We will use these precise descriptions of the local geometry to draw some conclusions of a global nature. First, we generalize the classical Gauss-Bonnet theorem.

Theorem 1.4. (Gauss-Bonnet Theorem) *Let V be a compact real analytic surface with isolated singularities p_1, \dots, p_R with an induced Riemannian metric on $V_0 = V \setminus \{p_1, \dots, p_R\}$ as above. Let K denote the Gauss curvature. According to Theorem 1.1, sufficiently small pointed neighborhoods of the p_i , $i = 1, \dots, R$, have connected components that are quasi-isometric to cones or horns. Let l_1, \dots, l_S be the leading coefficients in the length expansions (Theorem 1.3) for all the cone-like components.*

Then K is integrable over V_0 and

$$\chi(V) = R + \frac{1}{2\pi} \int_{V_0} K - \frac{1}{2\pi} \sum_{i=1}^S l_i,$$

where $\chi(V)$ is the Euler characteristic.

We now address some questions of analysis. The general problem here is the following: Given a Riemannian manifold V_0 which is the smooth part of a compact 'singular space' V , e.g. an analytic variety, how do the properties of the natural differential operators defined on V_0 differ from those in the smooth case (i.e. $V = V_0$)? Here the 'properties' can be of a very basic nature, like (i) below, or rather involved, like (iii). The Riemannian metric g induces a scalar product on differential forms. By $L^2(\wedge V_0)$ we denote the completion of the space of smooth forms on V_0 under this norm. The exterior derivative d and its transpose d^t act on elements of $L^2(\wedge V_0)$ in the sense of distributions.

Theorem 1.5. *Let (V, g) be as in Theorem 1.4.*

(i) *The L^2 Stokes theorem holds, i.e. if $\omega, \eta, d\omega, d^t\eta \in L^2(\wedge V_0)$ then*

$$(5) \quad (d\omega, \eta) = (\omega, d^t\eta).$$

(ii) *The Gauss-Bonnet operator $D_{GB} = d + d^t$ and the Laplace-Beltrami operator $\Delta = D_{GB}^2$ are self-adjoint as unbounded operators on $L^2(\wedge V_0)$ with domains $\mathcal{D}(D_{GB}) = H^1(\wedge V_0)$, $\mathcal{D}(\Delta) = H^2(\wedge V_0)$ (Sobolev spaces). Their spectra are discrete.*

(iii) *The L^2 -Euler characteristic of V , i.e. the index of the operator D_{GB} acting from even forms in $\mathcal{D}(D_{GB})$ to odd forms, equals*

$$\chi_{(2)}(V) = N + \frac{1}{2\pi} \int_{V_0} K - \frac{1}{2\pi} \sum_{i=1}^S l_i,$$

where N is the total number of cones and horns that occur near singularities of V .

(i) is equivalent to a property called 'uniqueness of ideal boundary conditions', see [17]. It says that no boundary terms occur in the integration by parts that is implicit in (5). It is conjectured to be true for all complex projective varieties with induced metrics, see [13] for related material. However, it is not true in general for compact real algebraic varieties, even if the singularities are isolated, see [17]. Note that (ii) does not say that Δ is essentially self-adjoint on $C_0^\infty(\wedge V_0)$. In fact, it need not be, as one can see for certain cones.

Outline of the proofs: The asserted quasi-isometry in Theorem 1.2 with any constant (resp. with the constant $1 + O(\varepsilon^\alpha)$) holds iff both of the following two statements are true (this is elementary; for a proof see Lemmas 3.1 and 3.2):

1. The angle β between the curves $\theta = \text{const}$ and the curves $r = \text{const}$ on K is bounded away from zero as $r \rightarrow 0$ (resp. tends to a right angle, with an error of order r^α).
2. The length $|\Phi_r|$ is bounded (resp. is $1 + O(r^\alpha)$).

We denote $\Phi_r = \partial\Phi/\partial r$ etc. For simplicity, we assume in this outline that $p = 0$ and g is induced from the Euclidean metric. Now $\beta = \angle(\Phi_r, \Phi_\theta)$. If we write $\Phi(r, \theta) = r\phi(r, \theta)$ with $|\phi| = 1$ then $\Phi_r = \phi + r\phi_r$ is a decomposition into a 'radial' component ϕ which is orthogonal to Φ_θ and a 'spherical' component $r\phi_r$. Therefore, the quasi-isometry is equivalent to the estimate $r|\phi_r| < C$ (resp. $r|\phi_r| < Cr^\alpha$).

Since the arc length parametrization is difficult to handle directly, we deal with arbitrary parametrizations $\Psi = r\psi, |\psi| = 1$, first and show (in Section 3) that the $r|\phi_r|$ -estimate follows from the analogous $r|\psi_r|$ -estimate plus the uniform boundedness, as $r \rightarrow 0$, of the integrals over K_r of the absolute geodesic curvature of K_r in K . The curvature estimate reduces to a length estimate in the unit tangent bundle, which can be proved by standard analytic techniques, see Section 4.

The problem is, then, to construct ψ satisfying $r|\psi_r| < Cr^\alpha$. This is easy in the model case of a family of curves given by hyperbolas, i.e. the family $x^a y^b = r$ in the (x, y) -plane, for positive integers a and b . Using resolution of singularities we can reduce the general case to this model. This will prove Theorem 1.2, see Section 2.

The asymptotic analysis of $l(r)$ (Section 5) is also most easily done on the blown-up space obtained by resolution of singularities. The curves K_r are then, locally, simply hyperbolas as above, but their length is measured with respect to a degenerating semi-Riemannian metric. In such a general situation, one always has an asymptotic expansion as stated, though the leading term may be $r^\gamma \log r$. We prove this using the Mellin transform, which reduces the problem to analytic continuation in the complex variable z of an integral of the form $\int_{\mathbb{R}^2} |q(x)|^{z+A} |p(x)|^{1/2} \chi(x) dx$ for real analytic functions p, q and $\chi \in C_0^\infty(\mathbb{R}^2)$, plus a decay estimate for large $\text{Im } z$. To do this, we adapt an idea by Bernshtejn and Gel'fand [2] and Atiyah [1], using resolution of singularities again. A separate consideration is needed to exclude a logarithm in the leading term, in our situation.

The Gauss-Bonnet Theorem is proved (in Section 6) by first excising small neighborhoods of the singular points and applying the classical Gauss-Bonnet Theorem to the resulting smooth manifold with boundary. The boundary integrals of geodesic curvature are interpreted as arc length variation. Here one needs the differentiated form of (the first term of) the asymptotics in Theorem 1.3. The integrability of K follows again by a standard argument, bounding $|K|$ by the area element of the unit normal bundle of V .

Theorem 1.5 follows easily from quasi-isometry invariance of some of the assertions and well-known results in the conic and horn-like cases. This is done in Section 7.

Further remarks and related work: All results are true for semi-analytic sets V as well, since the methods extend to this case. Clearly, the assumption $V \subset \mathbb{R}^N$ was only made for notational convenience. All that matters is that the metric on V_0 is locally induced by some local embedding. In any case, at least any smooth real analytic manifold can be embedded analytically in \mathbb{R}^N (see [19]), so probably the assumption $V \subset \mathbb{R}^N$ is no loss of generality.

In the preprint [5], L. Birbrair states a theorem that generalizes Theorem 1.1 to subanalytic surfaces with not necessarily isolated singularities, for g the Euclidean metric. However, there is no analogue of the sharper Theorem 1.2, and the parametrizations are not explicit. J.-M. Lion and J.-P. Rolin [22] prove a theorem on asymptotic expansions in a more general context which implies that the asymptotic series in Theorem 1.3 converges (i.e. $l(r) = l_1(r) + l_2(r) \log r$ with l_1, l_2 analytic in $r^{1/m}$); however, the statement about the leading term is not clear from [22]. The local geometry of semi-algebraic sets is also studied in [6]. In particular, the authors prove a Gauss-Bonnet Theorem for semi-algebraic sets. For more on curvature of sub-analytic sets see J. Fu [15].

If V is a *complex algebraic curve* then it falls in the class of surfaces considered here. In this case, Theorem 1.1 is much easier to prove. Questions of analysis have been treated in great detail by Nagase [23], Brüning-Lesch [9], [8] and Brüning-Peyerimhoff-Schröder [10].

The author wishes to thank M. Lesch for many helpful discussions and for suggesting to work on the spectral theoretic problems that motivated this study, and P. Milman for clarifying some questions on the resolution of singularities. Also, thanks to J. Fu and J. Tolksdorf for interesting discussions on the subject.

2. LOCAL METRIC STRUCTURE

In this section we carry out the main steps in the proof of Theorem 1.2.

We will use geodesic normal coordinates around p in \mathbb{R}^N ; these are given by the map

$$PC : [0, \infty) \times S^{N-1} \rightarrow \mathbb{R}^N$$

defined by $PC(r, \omega) = \gamma(r)$, where γ is the geodesic with $\gamma(0) = p$, $\gamma'(0) = \omega$. S^{N-1} denotes the unit sphere in $T_p \mathbb{R}^N$ with respect to $g|_p$, where we denote by g the metric on \mathbb{R}^N also. For r near zero, the form of the metric in normal coordinates is

$$(6) \quad PC^* g = dr^2 + r^2 g_{S^{N-1}}(r)$$

where $g_{S^{N-1}}(r)$ is a family of Riemannian metrics on S^{N-1} which depends smoothly on $r \geq 0$. We will denote this metric on S^{N-1} simply by $|\cdot|$.

We consider local parametrizations of K that satisfy (1), i.e. which are of the form

$$(7) \quad \begin{aligned} \Psi : (0, \varepsilon) \times (-A, A) &\rightarrow K, \\ (r, s) &\mapsto PC(r, \psi(r, s)) \end{aligned}$$

for some $A > 0$, $\varepsilon \leq \varepsilon_0$ and some map $\psi : (0, \varepsilon) \times (-A, A) \rightarrow S^{N-1}$.

The following proposition reduces the desired quasi-isometry statement to an estimate on derivatives of parametrizations Ψ , and to curvature estimates. It will be proved in the next section.

Proposition 2.1. *Suppose that there is a finite collection of parametrizations Ψ as in (7) whose ranges cover K and which satisfy the inequality, for some number C_0 :*

$$(8) \quad r|\psi_r| < C_0.$$

Also, suppose that there is a constant C_1 such that for the geodesic curvature κ of K_r in K we have

$$(9) \quad \int_{K_r} |\kappa| d\sigma < C_1$$

where $d\sigma$ denotes arc length.

Then there is a parametrization

$$\Phi : (0, \varepsilon) \times S^1 \rightarrow K$$

satisfying (3) and (2), with $\delta \leq C_0(6 + 4k(6 + 2C_1))$ where k is the number of Ψ needed to cover K .

In Section 4, we prove that (9) is satisfied in our situation, with C_1 only depending on V and g .

The following lemma shows that parametrizations Ψ satisfying a sharper estimate than (8) exist for certain model families of curves in \mathbb{R}^2 .

Lemma 2.2. *Let a, b be non-negative integers, not both zero, and let $A > 0$. There is a parametrization $\Psi(r, s) = (x(r, s), y(r, s))$ of each connected component of the family of curves*

$$x^a y^b = r, \quad r \in (0, 1), \quad |x| + |y| < A,$$

which satisfies

$$|\Psi_r| \leq r^{\frac{1}{a+b}-1}.$$

Here, $|\cdot|$ denotes the Euclidean metric on \mathbb{R}^2 .

Proof If $a = 0$ then we can take $\Psi(r, s) = (s, \pm r^{1/b})$, and the inequality is clearly satisfied. Similarly for $b = 0$.

Now assume $a > 0, b > 0$. We look at the curves in the first quadrant only, the other components are treated analogously. The desired parametrization $\Psi = (x, y)$ is defined by setting $s = x - y$. Namely, considering x, y as functions of r, s through the equations $s = x - y, r = x^a y^b$, with $x, y > 0$, one obtains by differentiating these equations with respect to r :

$$x_r = y_r = r^{-1} \left(\frac{a}{x} + \frac{b}{y} \right)^{-1}.$$

By the harmonic-geometric mean inequality, this is bounded above by

$$r^{-1}(a+b)^{-1}(x^a y^b)^{1/(a+b)} = (a+b)^{-1} r^{\frac{1}{a+b}-1}.$$

The lemma follows. \square

In order to reduce the general case to this model, we now 'blow up' \mathbb{R}^N to resolve the singularity p of V . That is, we find an analytic manifold X' and a proper analytic map $\pi : X' \rightarrow B$, where B is a small ball around p containing no other singularities of V , so that

- (a) π is a diffeomorphism $X' \setminus E \rightarrow B \setminus p$ where $E = \pi^{-1}(p)$.
- (b) The 'strict transform' $V' := \overline{\pi^{-1}(V \setminus p)}$ is a smooth surface.

Such an (embedded) resolution always exists, see [18] or [3]. For our application, we need to know some additional properties of this resolution:

Lemma 2.3. *The resolution $\pi : X' \rightarrow B$ can be chosen so that, in addition to (a) and (b) above, the following two properties are satisfied:*

- (c) *Near any point $q \in E \cap V'$ there are coordinates y_1, \dots, y_N centered at q such that, near q ,*

$$V' = \{y_3 = \dots = y_N = 0\} \quad \text{and}$$

$$r' := \pi^* r = \left| \prod_{i=1}^N y_i^{a_i} \right|$$

for some nonnegative integers a_i . Here r denotes the distance function to p on B .

- (d) *Let $h = r^{-2}(g - dr^2)$ be the semi-Riemannian metric on $B \setminus p$ obtained from the spherical part of g . Then there is a smooth semi-Riemannian metric h' on X' such that, on $X' \setminus E$, one has $h' = \pi^* h$.*

Remark Note that h does not extend continuously to $p \in B$. Thus the point of (d) is that after pulling back to X' , a smooth extension from $X' \setminus E$ to all of X' exists.

Before we prove this lemma, we finish the proof of Theorem 1.2. By Proposition 2.1 and Lemma 4.1, we are done if we construct parametrizations Ψ covering V which satisfy the estimate (8) with $C_0 = C\varepsilon^\alpha$. To do this, we first find parametrizations Ψ' of V' and then push them down to V : For each point $q \in V'_0 := V' \cap E$ let U_q be a neighborhood of q such that there are coordinates y_1, \dots, y_N on U_q as in (c) above. Below we construct, using Lemma 2.2, parametrizations Ψ'_q of the families of curves which make up the connected components of $V' \cap U_q \cap \{r' > 0\}$, such that Ψ'_q satisfies $r|\Psi'_{qr}|_{h'} < C_0 = C\varepsilon^\alpha$. Fix q and set $\Psi = \pi \circ \Psi'_q$. By construction, Ψ is of the form (7). We claim that ψ satisfies (8). Since $(PC^{-1}\Psi)_r = (1, \psi_r)$, we have $|\Psi_r|_h = |(PC^{-1}\Psi)_r|_{g_{SN-1}} = |\psi_r|_{g_{SN-1}}$. Therefore, we get $r|\psi_r|_{g_{SN-1}} = r|\Psi_r|_h = r|\Psi'_{qr}|_{h'} \leq C_0$ which was to be shown. Finally, since r' is proper, V'_0 is compact, so finitely many of the U_q cover V'_0 , and since r' must assume a positive minimum on the complement of their union U , there is an $\varepsilon_0 > 0$ such that $\{0 < r' < \varepsilon_0\}$ is contained in U . Thus, the assumptions of Proposition 2.1 are satisfied, so Theorem 1.2 is proved.

It remains to construct Ψ_q for each connected component of $V' \cap U_q \cap \{r' > 0\}$. If this set is non-empty then $a_3 = \dots = a_N = 0$. Then V' is parametrized

by y_1 and y_2 near q , and $r = y_1^{a_1} y_2^{a_2}$. We now use the parametrizations given by Lemma 2.2. The estimate (8) is then satisfied with respect to the Euclidean metric in the y_1, y_2 -plane and therefore also with respect to the smooth semi-Riemannian metric h' . Since the exponent of r in Lemma 2.2 is $\frac{1}{a+b} - 1$, the number C_0 can be chosen as $C\varepsilon^\alpha$ whenever $r < \varepsilon$ where C is a constant only depending on V (stemming from the length distortion between the Euclidean metric and the metric h' on U_q , i.e. $h' \leq Cg_{\text{eucl}}$ on U_q) and

$$\alpha = \min \frac{1}{a+b},$$

the minimum being taken over the finitely many coordinate neighborhoods U_q .

□

Proof of Lemma 2.3:

First, we need a few definitions: Let X be a real analytic manifold. A *normal crossings divisor (n.c.d.)* on X is a subset $E \subset X$ which, near any $q \in X$, is a union of coordinate hyperplanes, in a suitable local coordinate system near q . A set D has normal crossings with E if the coordinates can be chosen so that, in addition, D is a coordinate subspace. Finally, if f is a smooth function on X , we say that f has *product form with respect to E (and D)* if $\{f = 0\} \subset E$ and in one (and therefore any) such local coordinate system y_1, \dots, y_N near q with $y(q) = 0$, one has the representation

$$(10) \quad f(y) = h(y) \prod_{i=1}^N y_i^{a_i}$$

with h smooth and $h(0) \neq 0$, and non-negative integers a_i .

According to [3], a resolution can be obtained in the following way: Define a sequence

$$\begin{array}{ccccccc} X_M & \xrightarrow{\pi_{M-1}} & X_{M-1} & \rightarrow \dots \rightarrow & X_1 & \xrightarrow{\pi_0} & X_0 = B \\ E_M & & E_{M-1} & & E_1 & & E_0 = \emptyset \end{array}$$

of spaces X_j , n.c.d.s $E_j \subset X_j$, and proper maps $\pi_j : X_{j+1} \rightarrow X_j$ as follows: Let $X_0 = B, E_0 = \emptyset$. If X_j, E_j are defined, choose a suitable submanifold $D_j \subset X_j$ which is normal crossings with E_j , and let X_{j+1} be the elementary blow-up of X_j along D_j ; that is, essentially X_{j+1} is obtained from X_j by replacing D_j by its projectivized normal bundle, and π_j is the obvious projection map; also, set $E_{j+1} = \pi_j^{-1}(D_j \cup E_j)$, which is normal crossings again. It is one of the main theorems in [18] and [3] that the D_j can always be found such that after a finite number of such blowups, $X' := X_M$ is a resolution in the sense of (a) and (b) above, and such that V' and $E = E_M$ are normal crossings. Here, $\pi = \pi_{(M)}$ where $\pi_{(j)} = \pi_0 \circ \dots \circ \pi_{j-1}$. Also, one may choose D_j to lie in the singular set of $\pi_{(j)}^{-1}(V \setminus p)$ (resp. of V for $j = 0$).

Assume first that V is singular at p . We claim that (c) and (d) are satisfied for this resolution. The essential fact is that $V \cap B$ is only singular

at p , so that $\pi_0 : X_1 \rightarrow B$ is just the blow-up of p . If we identify B with a subset of \mathbb{R}^N via $x \equiv r\omega$ for $x = PC(r, \omega)$, then this can be described as $X_1 = (-\varepsilon_0, \varepsilon_0) \times S^{N-1} / \sim$ where $(t, \omega) \sim (-t, -\omega)$, and $\pi([t, \omega]) = t\omega$. Then $E_1 = \pi_0^{-1}(0) = \{t = 0\}$. To prove (c), we show inductively that $R_j := \pi_{(j)}^* r^2$ has product form with respect to E_j , for $j \geq 1$. This is clearly true for $j = 1$ since $\pi_0^* r^2 = t^2$ in the notation above. Then it follows directly from the definition of elementary blow-up that, if R_j has product form with respect to E_j and D_j has normal crossings with E_j , then $R_{j+1} = \pi_j^* R_j$ has product form with respect to E_{j+1} . (Use projective coordinates near points in $\pi_j^{-1}(D_j)$.) Therefore, $R_M = \pi^* r^2$ has the form (10). Since $R_M \geq 0$, h is positive and all a_i are even. Since $E = \{R_M = 0\}$, one of the a_i is non-zero for $q \in E$, so one may modify the coordinates such that $h \equiv 1$. This proves (c).

To prove (d), we note that the pull-back metric on $(-\varepsilon_0, \varepsilon_0) \times S^{N-1}$ is $dt^2 + t^2 g_{S^{N-1}}(t)$ with $g_{S^{N-1}}(t)$ as in (6) for $t \geq 0$ and $g_{S^{N-1}}(-t) = A^* g_{S^{N-1}}(t)$ where A is the antipodal map. This shows that $h_1 := g_{S^{N-1}}(t)$ is well-defined and smooth on X_1 , and satisfies $h_1 = \pi_0^* h$ on $X_1 \setminus E_1$. Therefore, $h' = (\pi_1 \circ \dots \circ \pi_{r-1})^* h_1$ is also smooth.

Finally, if V is smooth at p then just blow up p once, then (c) and (d) are satisfied by the same argument. \square

3. BASICS ON QUASI-ISOMETRY

Lemma 3.1. *Let Ψ be a local parametrization as in (7). Then*

$$(11) \quad \Psi^* g \cong dr^2 + H(r, s)^2 ds^2$$

for some function H if and only if

$$r|\psi_r(r, s)| < C$$

for some constant C . In this case, one can take $H = r|\psi_s|$. Then the quasi-isometry constant depends only on C , and for $C < 1/2$ can be taken to be $1 + 2C$.

Proof : From (6) and $\Psi = PC \circ (r, \psi)$ we have

$$\begin{aligned} \Psi^* g &= dr^2 + r^2 |d\psi|^2 \\ &= dr^2 + r^2 |\psi_r dr + \psi_s ds|^2 \\ &= (1 + |A|^2) dr^2 + 2A \cdot B dr ds + |B|^2 ds^2 \end{aligned}$$

where $A = r\psi_r, B = r\psi_s$. We now use the following elementary lemma.

Lemma 3.2. *Let $g = a dx^2 + 2b dx dy + c dy^2$ be a positive definite quadratic form, and let $\tilde{g} = a dx^2 + c dy^2$ be its 'diagonal part'. Set*

$$T = \frac{|b|}{\sqrt{ac}}.$$

Then one has

$$(1 - T)\tilde{g} \leq g \leq (1 + T)\tilde{g},$$

and these inequalities are sharp.

Remark: Note that T is the cosine of the angle between the x -axis and the y -axis with respect to the Riemannian metric g on \mathbb{R}^2 . Therefore, if a, b, c are allowed to depend on x and y , one obtains that $g \cong \tilde{g}$ if and only if this angle is bounded away from zero, and the quasi-isometry constant is close to one if this angle is close to $\pi/2$.

Proof The left inequality is equivalent to $Ta dx^2 + 2b dx dy + Tc dy^2 \geq 0$, and this is equivalent to $Ta \geq 0$ and $Ta \cdot Tc - b^2 \geq 0$. Similarly for the right inequality. This implies the claim. Note that always $T < 1$ since $g > 0$ implies $ac - b^2 > 0$. \square

Here we have, for fixed (r, s) , $a = 1 + |A|^2$, $b = A \cdot B$ and $c = |B|^2$, so

$$T = \frac{|A \cdot B|}{|B|\sqrt{1+|A|^2}} \leq \frac{|A|}{\sqrt{1+|A|^2}}.$$

Also, one clearly has $dr^2 + |B|^2 ds^2 \approx_{1+|A|^2} (1+|A|^2)dr^2 + |B|^2 ds^2$. Therefore, from Lemma 3.2 we get a quasi-isometry factor, for fixed (r, s) , of at most $(1+|A|^2)(1 - \frac{|A|}{\sqrt{1+|A|^2}})^{-1}$. An easy calculation shows that this is bounded for A bounded, and for $|A| < 1/2$ is bounded above by $1 + 2A$. Finally, looking at the length of $\partial/\partial r$, one sees that $|A| < C$ is also necessary for the asserted quasi-isometry.

\square

The following lemma relates arbitrary parametrizations to (unrenormalized) arc length parametrizations.

Lemma 3.3. *Let $\Omega(r, t) = PC(r, \omega(r, t))$, $(r, t) \in (r_1, r_2) \times (t_1, t_2)$ be a local parametrization of K with ω arc length on the sphere, i.e. $|\omega_t| \equiv 1$. Assume that there is a parametrization $\Psi : (0, \varepsilon) \times (-A, A) \rightarrow K$ whose range contains the range of Ω and which satisfies the inequality (8), and that (9) holds.*

Then, if $r|\omega_r(r, t)| < C$ holds for one value $t = t_0$, it holds for all t (with C replaced by $C + C_0 C_2$ where $C_2 = 6 + 2C_1$).

Proof We may assume that Ψ and Ω trace the lines $r = \text{const}$ in the same direction. Define the function $b(r)$ by the equation

$$\omega(r, t_0) = \psi(r, b(r)).$$

Then, if we set $l(r, s) = \int_{b(r)}^s |\psi_s(r, \sigma)| d\sigma$, we must have

$$\omega(r, t_0 + l(r, s)) = \psi(r, s).$$

Differentiating in r , we get $\omega_r + l_r \omega_t = \psi_r$, evaluated at corresponding points. From condition (8) and $|\omega_t| = 1$ we get

$$(12) \quad r|\omega_r(r, t_0 + l(r, s))| < C \Leftrightarrow r|l_r(r, s)| < C,$$

where C must be replaced by $C + C_0$ in either direction of the implication. Now we have $l_r = a(r) + \int_{b(r)}^s \frac{\partial}{\partial r} |\psi_s| ds$ where $a(r) = -b'(r)|\psi_s(r, b(r))|$. The

variation of length formula says that

$$\int_{b(r)}^s \frac{\partial}{\partial r} |\psi_s| ds = \left(\frac{\psi_s}{|\psi_s|}, \psi_r \right) \Big|_{b(r)}^s - \int_{b(r)}^s (N, \psi_r) \kappa d\sigma$$

where κ is the geodesic curvature, N is the unit normal, and σ denotes arc length. From (8) and (9) we get $r|l_r - a| \leq C_0(2 + C_1)$. Therefore, the right side of (12) is equivalent to $r|a(r)| < C$, where the constant gets worse by $C_0(2 + C_1)$. Now since this last condition is independent of s , and since the left side of (12) is true for $s = b(r)$ by assumption, it must be true for all s . Altogether we lose at most $C_0(6 + 2C_1)$ in the constant.

□

Proof of Proposition 2.1

First we consider a parametrization

$$\begin{aligned} \Omega : (0, \varepsilon) \times [0, \infty) &\rightarrow K \\ (r, t) &\mapsto PC(r, \omega(r, t)) \end{aligned}$$

for which ω is arc length on the sphere, i.e. $|\omega_t| \equiv 1$. Such a parametrization is determined when the 'initial' curve $r \mapsto \Omega(r, 0)$ and an orientation for the family of curves K_r are prescribed. We assume such an orientation given and prescribe the initial curve by choosing any one local parametrization ψ and setting

$$(13) \quad \omega(r, 0) = \psi(r, 0).$$

Note that $|\Omega_t| = r$ and $\text{length}(K_r) = l(r)$ imply that

$$(14) \quad \omega(r, \tilde{l}(r)) = \omega(r, 0)$$

for $\tilde{l}(r) = l(r)/r$. We now prove that the parametrization ω thus defined satisfies the inequality

$$(15) \quad r|\omega_r(r, t)| < C$$

for $t \leq \tilde{l}(r)$, with $C = (1 + kC_2)C_0$. Fix $r_0 \in (0, \varepsilon)$. We prove (15) for $r = r_0$, but with C independent of r_0 : Since each of the finitely many given parametrizations Ψ covers an interval on the curve $\gamma = K_{r_0}$, one can choose points $p_0 = \Omega(r_0, 0), p_1, p_2, \dots, p_k = p_0$ in this order on γ such that each subarc from p_i to p_{i+1} of γ is contained in the range of some parametrization Ψ_i satisfying the estimates stated in the proposition. Define $0 = t_0 < t_1 < \dots < t_k = \tilde{l}(r_0)$ by letting rt_i (for $i < k$) be the distance from p_0 to p_i measured along γ in the positive direction; then $\Omega(r_0, t_i) = p_i$. We prove by induction on i that (15) is true for $t \leq t_i$. For $i = 0$, $t_0 = 0$ and therefore (15) is true by (13), with $C = C_0$. Suppose it is true for $t \leq t_i$ with some constant $C = C_{(i)}$. Since the range of Ψ_i is open, it will contain the range of Ω when restricted to a small interval around r_0 times some interval containing $[t_i, t_{i+1}]$. Thus Lemma 3.3 implies that (15) holds for $t \in [t_i, t_{i+1}]$, with $C = C_{(i+1)} = C_{(i)} + C_2C_0$. Altogether, we get that (15) holds for all $t \leq \tilde{l}(r)$ with $C = (1 + kC_2)C_0$.

Now we renormalize ω by setting

$$(16) \quad \phi(r, \theta) = \omega(r, \theta \tilde{l}(r)/2\pi).$$

We show that ϕ also satisfies the inequality

$$(17) \quad r|\phi_r(r, \theta)| < C,$$

where now $C = (3 + 2kC_2)C_0$ and $\theta \in [0, 2\pi]$. Differentiating (14) in r , we get $\omega_r(r, 0) = \omega_r(r, \tilde{l}(r)) + \tilde{l}'(r)\omega_t(r, \tilde{l}(r))$. Now (15) and $|\omega_t| = 1$ give $r|\tilde{l}'| < (2 + kC_2)C_0$, and by (16) this in turn implies (17).

Now set $\Phi = PC \circ (r, \phi)$. By definition, we have $|\Phi_\theta| = r|\phi_\theta| = r\tilde{l}(r)/2\pi = l(r)/2\pi$, and Lemma 3.1, applied to Φ , gives (2). \square

4. PROOF OF THE CURVATURE BOUND (9)

Lemma 4.1. *Condition (9) is satisfied, with C_1 only depending on V and g .*

Proof Let NK_r denote the unit normal bundle of K_r in V , a one-dimensional submanifold of the unit tangent bundle of V_0 . It is an elementary fact from differential geometry that

$$\int_{K_r} |\kappa| \leq \text{length}(NK_r).$$

So it remains to show that these lengths are uniformly bounded as $r \rightarrow 0$. Set $U = \bigcup_{r \in (0, \varepsilon)} NK_r$. Now U is a semi-analytic subset of $T^1\mathbb{R}^N$, the unit tangent bundle of (\mathbb{R}^N, g) . This can be seen as follows: Assume V is given as $\{f_1 = \dots = f_L = 0\}$ near p . If V is a complete intersection at p , i.e. $L = N - 2$, then we can write

$$U = \{(x, v) \in T^1\mathbb{R}^N : df_{i|_x}(v) = 0, \omega_x(v) = 0, x \in K \setminus p\};$$

here the analytic one-form ω is defined as $\omega = *(d(r^2) \wedge df_1 \wedge \dots \wedge df_L)$ where $*$ denotes the Hodge star operator. If V is not a complete intersection at p then define, for every subset $S \subset \{1, \dots, L\}$ with $N - 2$ elements, the one-form $\omega_S = *(d(r^2) \wedge \bigwedge_{i \in S} df_i)$ and use the conjunction of $\omega_{S|_x}(v) = 0$ over all S in the description of U .

We now have the following situation: $X = T^1\mathbb{R}^N$ is a Riemannian real analytic manifold, with metric induced from the metric on \mathbb{R}^N . $U \subset X$ is a two-dimensional semi-analytic subset and $R := r^2 : X \rightarrow \mathbb{R}$ is a proper analytic function which is non-constant on U . We claim that this implies that the level sets of R have bounded lengths (i.e. one-dimensional Hausdorff measure), for bounded R . This can be proved as follows (see [4] for definitions and theorems used): \overline{U} is semianalytic and two-dimensional. Let $\pi : Y \rightarrow X$ be a uniformization of \overline{U} ; this means that Y is smooth and two-dimensional, π is proper, and $\pi(Y) = \overline{U}$. (A uniformization can be obtained from a resolution, but also in a simpler way, see [4]). Let h be the pull-back semi-Riemannian metric on Y . Now $R' = \pi^*R : Y \rightarrow \mathbb{R}$ is proper. Clearly, the length of any level line $\{R = t\}$ in X is bounded by the length (with

respect to h) of the level line $\{R' = t\}$ in Y . Therefore, we are now in the smooth two-dimensional case, which is easy. For example, one can blow-up (fairly explicitly) Y to put R' into local product form, then the level sets are families of hyperbolas locally, and since the pull-back semi-metric is smooth, they have uniformly bounded length (compare Section 5). \square

5. ASYMPTOTICS OF $l(r)$

We investigate $l(r)$ by evaluating it on the blown-up space X' , described in Section 2. The pull-back metric then degenerates on the preimage of $\{r = 0\}$.

We first analyze the model case, where the curves are given in \mathbb{R}^2 by $x^a y^b = r$, but the metric may degenerate on the coordinate axes:

Proposition 5.1. *Let a, b be nonnegative integers, $a + b > 0$. Let h be a real analytic semi-Riemannian metric defined near zero in \mathbb{R}^2 which is Riemannian for $x^a y^b \neq 0$, and $\chi \in C_0^\infty(\mathbb{R}^2)$ a cutoff function which is non-negative and equals one near the origin. Let $l_\chi(r) = l_{\chi, h}(r)$ be the length of the curve $x^a y^b = r$, weighted by χ , i.e.*

$$l_\chi(r) = \int |\psi_s(r, s)|_{h_{\psi(r, s)}} \chi(\psi(r, s)) ds$$

for some parametrization ψ of this family of curves.

Then $l_\chi(r)$ has an asymptotic expansion, as $r \rightarrow 0$

$$l_\chi(r) \sim \sum_{i, j} C_{i, j} r^i (\log r)^j$$

where i ranges over $\frac{1}{m}\mathbb{N}_0$ for some fixed $m \in \mathbb{N}$, and $j \in \{0, 1\}$. This expansion may be differentiated (arbitrarily often) term by term.

If, in addition, h is the pull-back of a non-degenerate (i.e. Riemannian) metric under an analytic map which has injective differential for $x^a y^b \neq 0$, then the leading term is of the form r^γ .

Note The following example shows that, without further assumptions, $\log r$ may appear in the leading term: Let $h = y^2 dx^2 + x^2 dy^2$ and $a = b = 1$. Then, parametrizing by $s = x$, one has

$$|\psi_s|_h = |(1, -r/x^2)|_h = ((r/x)^2 + x^2(-r/x^2))^{1/2} = \sqrt{2} r/x$$

which is easily seen to imply $l_\chi(r) \sim -\sqrt{2} r \log r$.

Proof Clearly, l_χ is bounded, smooth for $r > 0$, and has compact support. We use the Mellin transform, defined, for a bounded compactly supported function l , by

$$(Ml)(z) = \int_0^\infty r^z l(r) \frac{dr}{r}$$

with $z \in \mathbb{C}$. This integral is defined and holomorphic for $\operatorname{Re}(z) > 0$. As is well-known (see [20], for example), an asymptotic expansion for l as $r \rightarrow 0$

(with derivatives) is equivalent to the existence of a meromorphic continuation of $Ml(z)$ to the whole z -plane, with a pole of order μ at z_0 corresponding to a term $r^{-z_0}(\log r)^{\mu-1}$ in the asymptotics, and such that one has decay for large imaginary part, i.e.

$$(18) \quad |Ml(z)| \leq C_N |\operatorname{Im} z|^{-N}$$

for any N , uniformly for bounded $\operatorname{Re} z$ and away from the poles. Therefore, we have to show the existence of such a continuation with at most double poles, at points in $-\mathbb{N}_0/m$. The additional statement on the leading term will be proved directly.

We assume $a > 0, b > 0$, the other case is only slightly different, but easier.

A simple calculation, for example using the parameter $s = x$, shows that, as measures,

$$|\psi_s|_{h_\psi} ds \frac{dr}{r} = |(bx, -ay)|_{h_{(x,y)}} \frac{dx}{x} \frac{dy}{y}.$$

Therefore, we get

$$(19) \quad Ml_\chi(z) = \int_0^\infty \int_0^\infty x^{az} y^{bz} p(x, y)^{1/2} \chi(x, y) \frac{dx}{x} \frac{dy}{y}$$

where we have used $r = x^a y^b$, and where the analytic function p is defined as $p(x, y) = |(bx, -ay)|_{h_{(x,y)}}^2$. If we set $x = x_1^2, y = x_2^2$ then (19) becomes

$$Ml_\chi(z) = \int_{-\infty}^\infty \int_{-\infty}^\infty |x_1|^{2az-1} |x_2|^{2bz-1} p(x_1^2, x_2^2)^{1/2} \chi(x_1^2, x_2^2) dx_1 dx_2,$$

and the asymptotics follows from the following proposition:

Proposition 5.2. *Let p, q be real analytic functions defined in a neighborhood U of the support of a function $\rho \in C_0^\infty(\mathbb{R}^n)$. Let $A, B \in \mathbb{R}$, $B \geq 0$. For $z \in \mathbb{C}$, $\operatorname{Re} z \geq -A$, define*

$$f(z) = \int_{\mathbb{R}^n} |q(x)|^{z+A} |p(x)|^B \rho(x) dx.$$

Then

- (a) *f can be extended meromorphically to all of \mathbb{C} , with poles of order at most n .*
- (b) *If, for every $x \in \operatorname{supp} \rho$, $q(x) \neq 0$ implies $p(x) \neq 0$ and $dq|_x \neq 0$ then*

$$|f(z)| \leq C_N |\operatorname{Im} z|^{-N}$$

for any N , uniformly in $|\operatorname{Im} z| \geq 1$ and for bounded $\operatorname{Re} z$.

Proof Part (a) is a consequence of a more general theorem by Gel'fand and Bernshtejn, see [2]. We sketch their proof and show how one obtains (b). First, we use the resolution of singularities theorem to find an analytic manifold X and an analytic proper map $\pi : X \rightarrow U$ so that (1) π is a diffeomorphism outside $\{pq = 0\}$ and (2) the pull-back $\pi^*(pq)$ has local product form; i.e. near any point in X there are analytic local coordinates

(y_1, \dots, y_n) centered at that point such that $\pi^*(pq)$ is a monomial in the y_i times a non-vanishing analytic function. Then π^*p and π^*q must have local product form, too, and if we write $\pi^*(dx_1 \wedge \dots \wedge dx_n) = S(y)dy_1 \wedge \dots \wedge dy_n$, then S must have local product form (this follows from Hilbert's Nullstellensatz – resp. Rückert's Nullstellensatz in the analytic case, see [24] – since S vanishes only where $\pi^*(pq)$ vanishes, even after local complexification). All this implies that we can use a partition of unity on X to write f as a finite sum of terms of the form

$$g(z) = \int_{\mathbb{R}^n} |\tilde{q}(y)|^{z+A} |\tilde{p}(y)|^B |S(y)| \tilde{\rho}(y) dy$$

where $\tilde{\rho} \in C_0^\infty(\mathbb{R}^n)$, $\tilde{q}(y) = \pi^*q(y) = y^\alpha h_1(y)$, $\tilde{p}(y) = \pi^*p(y) = y^\beta h_2(y)$, $S(y) = y^\gamma h_3(y)$ with multi-indices α, β, γ and h_1, h_2, h_3 non-vanishing on $\text{supp } \tilde{\rho}$. For each g we have one of the following two cases:

(I) $\alpha \neq 0$, i.e. $\tilde{q}(0) = 0$.

(II) $\alpha = 0$; then $\gamma = 0$ by (1) above, and under the additional condition in part (b) of the proposition also $\beta = 0$, $d\tilde{q} \neq 0$.

We first consider case (II): If $\alpha = 0$ then $|\tilde{q}|^z$ and therefore g is entire in z , which proves (a). For (b), we may assume w.l.o.g. that $\tilde{q}(y) = e^{y_1}$, then

$$g(R + iI) = \int_{-\infty}^{\infty} e^{iIy_1} \left(e^{(R+A)y_1} \int_{\mathbb{R}^{n-1}} |\tilde{p}|^B |S| \tilde{\rho} dy' \right) dy_1.$$

Since $\tilde{p}, S \neq 0$ on $\text{supp } \tilde{\rho}$, this is the Fourier transform of a smooth compactly supported function, therefore rapidly decreasing in I .

We now consider case (I): By a smooth change of coordinates, we may arrange $h_1 \equiv 1$. Then (with $\delta = \alpha A + \beta B + \gamma$)

$$g(z) = \int_{\mathbb{R}^n} |y_1|^{\alpha_1 z + \delta_1} \dots |y_n|^{\alpha_n z + \delta_n} \chi(y) dy, \quad \chi \in C_0^\infty(\mathbb{R}^n).$$

The following stronger result now implies the proposition, with m equal to the least common multiple of all the *positive* a_i , over all i and g : If $\chi \in C_0^\infty(\mathbb{R}^n)$ then the function

$$g(z_1, \dots, z_n) = \int_{\mathbb{R}^n} |y_1|^{z_1} \dots |y_n|^{z_n} \chi(y) dy$$

has a meromorphic continuation from $\{\text{Re } z_i \geq 0 \forall i\}$ to \mathbb{C}^n , with at most simple poles on the hyperplanes $\{z_i = -s\}, s \in \mathbb{N}$, and on $\{(z_1, \dots, z_n) : \text{dist}(z_i, -\mathbb{N}) > \varepsilon, |\text{Re } z_i| < \varepsilon^{-1} \forall i\}$ one has for any N

$$|g(z_1, \dots, z_n)| \leq C_{\varepsilon, N} \prod_{i=1}^n |\text{Im } z_i|^{-N}$$

where C_N is bounded in terms of the C^M norm of χ for some $M = M(N)$. This claim follows for $n = 1$ directly from standard facts about distributions, see [16], and for $n > 1$ by induction. \square

It remains to prove the additional statement in Proposition 5.1 about the leading term. Assume $h = \pi^*g$ for an analytic map $\pi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^N, 0)$

as in the proposition and a Riemannian metric g on \mathbb{R}^N (everything defined near zero). Then $|\psi_s|_h = |\partial_s(\pi \circ \psi)|_g$. Assume $b > 0$. We will use the parametrization by $s = x$ now, i.e.

$$\psi(r, s) = (s, r^{1/b}/s^{a/b}).$$

A short calculation (as above) shows that

$$(20) \quad \partial_s(\pi(\psi(r, s))) = \frac{1}{bx}(bx\pi_x - ay\pi_y) \in \mathbb{R}^N$$

where $(x, y) = \psi(r, s)$. Now the essential fact is the following: As function of r and s , every component of (20) is a Laurent series in $s^{1/b}$ whose coefficients are power series in $r^{1/b}$. Since it is a derivative in s , the s -residue, i.e. the coefficient of s^{-1} , vanishes. This means that no component of the analytic function $bx\pi_x - ay\pi_y$ contains a monomial of the form $(x^a y^b)^\alpha$. Also, this function does not vanish identically by the non-degeneracy hypothesis on π . In particular, letting $\gamma \in \mathbb{Q}$ be the largest number such that $(x^a y^b)^\gamma$ divides $bx\pi_x - ay\pi_y$, we have

$$bx\pi_x - ay\pi_y = (x^a y^b)^\gamma Q(x, y)$$

with $Q : \mathbb{R}^2 \rightarrow \mathbb{R}^N$ analytic in certain fractional powers of x and y and $Q(0, 0) = 0$, and such that Q does not vanish identically on $\{x = 0\} \cup \{y = 0\}$. Since g is nondegenerate, $q = |Q|_g^2$ has the same properties.

To finish the proof, it is clearly enough to show that, for some constants $0 < c < C < \infty$, we have

$$cr^\gamma < l_\chi(r) < Cr^\gamma,$$

for small r . From (20) and the definition of q , we have

$$l_\chi(r) = r^\gamma I(r)/b \text{ where } I(r) = \int q(x, y)^{1/2} \chi(x, y) \frac{dx}{x}$$

where always $y = r^{1/b}/x^{a/b}$. To show $I < C$, we observe that $q(0, 0) = 0$ implies $q(x, y) \leq x^B + y^B$ for some $B > 0$. Since $\frac{dx}{x} = \frac{b}{a} \frac{dy}{y}$, we have $I \leq \int_0^1 x^B dx/x + b/a \int_0^1 y^B dy/y < C$.

To show $I > c$, we recall that q does not vanish identically on at least one coordinate axis, say the x -axis. Then, $q^{1/2} > c_0 > 0$ on some rectangle $J \times [0, \delta]$ on which $\chi = 1$, where J is an open interval on the x -axis and $\delta > 0$. Then $I \geq c_0 \int_J dx/x =: c$ whenever r is so small that $x^a y^b = r, x \in J$ implies $y < \delta$. This finishes the proof of the proposition.

□

Proof of Theorem 1.3: Apply the proposition to the semi-Riemannian metric $h_1 = \pi^*g$ and to $h_2 = h'$ from Lemma 2.3 (d), and use a partition of unity on X' . Clearly $l_{\chi, h_1} = r l_{\chi, h_2}$. Therefore, the leading term has exponent $\gamma \geq 1$. Also, since g is non-degenerate, there is no logarithm. □

6. THE GAUSS-BONNET THEOREM

Here we prove Theorem 1.4. The integrability of K , i.e. the finiteness of $\int_{V_0} |K|$, is proved in a similar way as the bound on geodesic curvature (9), only simpler. We sketch the argument in three dimensions; for details see [14]: From the fact that K is the Jacobian of the differential of the Gauss map one easily sees that for any open, relatively compact set $U \subset V_0$ one has $\int_U |K| \leq \text{area}(NU)$, where NU is the unit normal bundle of U , with the induced metric. Since V_0 is semi-analytic, so is NV_0 , so it has finite area by compactness of V . Therefore, $\int_{V_0} |K|$ is finite also.

To prove the Gauss-Bonnet formula, we choose $\varepsilon_0 > 0$ small enough so that Theorem 1.2 applies for each p_i (but here we only need the topological statement), and let $U_\varepsilon = \bigcup_{i=1}^R B_\varepsilon(p_i)$ and $V_\varepsilon = V \setminus U_\varepsilon$. For $\varepsilon < \varepsilon' < \varepsilon_0$ use the additivity of the Euler characteristic and $V = V_\varepsilon \cup U_{\varepsilon'}$ to obtain

$$(21) \quad \chi(V) = \chi(V_\varepsilon) + \chi(U_{\varepsilon'}) - \chi(V_\varepsilon \cap U_{\varepsilon'}).$$

Now $U_{\varepsilon'}$ is homotopic to $\{p_1, \dots, p_R\}$, so

$$\chi(U_{\varepsilon'}) = R,$$

and $V_\varepsilon \cap U_{\varepsilon'}$ is homotopic to a disjoint union of circles, so

$$\chi(V_\varepsilon \cap U_{\varepsilon'}) = 0.$$

Applying the classical Gauss-Bonnet Theorem (see [25], Ch. 6) to the manifold with boundary V_ε , we get

$$(22) \quad \chi(V_\varepsilon) = \frac{1}{2\pi} \int_{V_\varepsilon} K + \frac{1}{2\pi} \int_{\partial V_\varepsilon} \kappa,$$

where κ is the geodesic curvature of $\partial V_\varepsilon = \partial U_\varepsilon$ with respect to the normal pointing outward from V_ε . Now the variation of arc length is geodesic curvature (see [25], Ch. 9), i.e. (with the previous orientation)

$$\frac{d}{d\varepsilon} l(\partial V_\varepsilon) = - \int_{\partial V_\varepsilon} (\partial/\partial r, N) \kappa$$

where N is the unit normal (pointing away from the p_i) and $\partial/\partial r$ is taken with respect to the coordinates described in Theorem 1.2. Now this theorem implies that the scalar product $(\partial/\partial r, N)$ tends to 1 as $r = \varepsilon \rightarrow 0$, uniformly in θ . Since $\int_{V_\varepsilon} |\kappa|$ is uniformly bounded (Lemma 4.1), we get

$$\frac{d}{d\varepsilon} l(\partial V_\varepsilon) = - \int_{\partial V_\varepsilon} \kappa + o(\varepsilon)$$

By Theorem 1.3, we have

$$\frac{d}{d\varepsilon} l(\partial V_\varepsilon)|_{\varepsilon=0} = \sum_{i=1}^S l_i.$$

Therefore, taking the limit $\varepsilon \rightarrow 0$ in (21) and (22) we get the result. \square

7. ANALYSIS NEAR A SINGULAR POINT

Here we prove Theorem 1.5. For a differential operator D on forms, denote by D_{\min} and D_{\max} the unbounded operators in $L^2 := L^2(\wedge V_0)$ with domains $\{\omega \in L^2 : \exists \omega_n \in C_0^\infty(V_0), \omega_n \rightarrow \omega \text{ and } D\omega_n \rightarrow D\omega \text{ in } L^2\}$ and $\{\omega \in L^2 : D\omega \in L^2\}$, respectively.

Proof of (i): First, we show that validity of the L^2 Stokes Theorem depends only on the quasi-isometry class of the metric. This is not obvious, since the local expression for d^t contains derivatives of the metric coefficients. But it follows directly from the fact that (5) is equivalent to the statement

$$d_{\min} = d_{\max}$$

and the fact that these two operators (in particular their domains) are quasi-isometry invariants. To see this equivalence, note that (5) is equivalent to

$$(23) \quad (d_{\max})^* = (d^t)_{\max}$$

where the star denotes the functional analytic adjoint. Also, it is easy to see that $(d_{\min})^* = (d^t)_{\max}$. Since d_{\min} and d_{\max} are closed operators, the claim follows. For details see [17].

By Theorem 1.1 it is therefore enough to prove (5) for metrics on V_0 that are exactly horn- or cone-like near the singularities. This is a special case of a well-known result by Cheeger [11], Theorem 2.2.

Proof of (ii): By (i) and (23), the operator $d_{\max} + (d^t)_{\max}$ is self-adjoint. Its domain is $\{\omega \in L^2 : d\omega, d^t\omega \in L^2\} = \mathcal{D}(D_{\text{GB},\max}) =: H^1(\wedge V_0)$, where the first equality follows from the orthogonality of the ranges of d and d^t . Since $H^2(\wedge V_0) = \{\omega \in L^2 : d\omega, \Delta\omega \in L^2\} = \mathcal{D}((D_{\text{GB},\max})^2)$, one also has self-adjointness of Δ . Finally, discreteness of the spectrum is a quasi-isometry invariant by [7], Lemma 2.17, and was proved for cones and horns by Cheeger [12] and Lesch-Peyerimhoff [21].

Proof of (iii): One can argue as in the proof of Theorem 1.4: $\chi_{(2)}$ is also additive. For manifolds with boundary (and metrics smooth up to the boundary) or their open interior, one has $\chi_{(2)} = \chi$, so only the term $\chi_{(2)}(U_{\varepsilon'})$ in (21) needs to be reevaluated. Now $\chi_{(2)}$ is a quasi-isometry invariant, and on cones and horns it equals one by [11], Lemma 3.4. Since $\chi_{(2)}$ is calculated on the smooth part V_0 , one has $\chi_{(2)}(U_{\varepsilon'}) = \chi_{(2)}(U_{\varepsilon'} \setminus p_1, \dots, p_R) = N$. \square

REFERENCES

- [1] M. ATIYAH, *Resolution of singularities and division of distributions*, Comm. Pure Appl. Math. **23**(1970), 145-150.
- [2] I. N. BERNSHTEIN AND S. I. GEL'FAND, *Meromorphic property of the functions P^λ* (English. Russian original) Funct. Anal. Appl. **3**(1969), 68-69; translation from Funkts. Anal. Prilozh. **3**(1969), 84-85.
- [3] E. BIERSTONE AND P. D. MILMAN, *Canonical desingularization in characteristic zero by blowing up the maximum strata of a local invariant*, Inv. Math. **128**(1997), 207-302.
- [4] E. BIERSTONE AND P. D. MILMAN, *Semianalytic and subanalytic sets*, IHES Publ. Math. **67** (1988), 5-42.

- [5] L. BIRBRAIR, *Local bi-Lipschitz classification of 2-dimensional semialgebraic sets*, Preprint 1998.
- [6] L. BRÖCKER AND M. KUPPE AND W. SCHEUFLER, *Inner metric properties of 2-dimensional semi-algebraic sets*, in: Real algebraic and analytic geometry (Segovia 1995), Rev. Mat. Univ. Complut. Madrid **10**(1997), 51-78.
- [7] J. BRÜNING AND M. LESCH, *Hilbert complexes*, J. Funct. Anal. **108** (1992), 88–132.
- [8] J. BRÜNING AND M. LESCH, *Kähler–Hodge theory for conformal complex cones*, Geom. Funct. Anal. **3** (1993), 439–473.
- [9] J. BRÜNING AND M. LESCH, *On the spectral geometry of algebraic curves*, J. reine angew. Math. **474** (1996), 25–66.
- [10] J. BRÜNING, N. PEYERIMHOFF, AND H. SCHRÖDER, *The $\bar{\partial}$ -operator on algebraic curves*, Commun. Math. Phys. **129** (1990), 525–534.
- [11] J. CHEEGER, *On the Hodge theory of Riemannian pseudomanifolds*, Proc. Symp. Pure Math. **36** (1980), 91–146.
- [12] J. CHEEGER, *Spectral geometry of singular Riemannian spaces*, J. Diff. Geom. **18** (1983), 575–657.
- [13] J. CHEEGER, M. GORESKY, AND R. MACPHERSON, *L^2 -cohomology and intersection homology of algebraic varieties*, Annals of Math. Studies **102** (1982), 303–340.
- [14] G. EULERING, *Integrale Krümmungskonstanten algebraischer Untermannigfaltigkeiten von Räumen konstanter Krümmung*, Doctoral Thesis, Münster 1995.
- [15] J. FU, *Curvature measures of subanalytic sets*, Am. J. Math. **116** (1994), 819–880.
- [16] I. M. GEL'FAND AND G. E. SHILOV, *Generalized Functions, Vol. I*, Academic Press, New York, 1964.
- [17] D. GRIESER AND M. LESCH, *On the L^2 Stokes theorem and Hodge theory for singular algebraic varieties*, Preprint 1998.
- [18] H. HIRONAKA, *Resolution of singularities of an algebraic variety over a field of characteristic zero*, Ann. of Math. **79** (1964), 109–326.
- [19] M. W. HIRSCH, *Differential Topology*, Springer 1976.
- [20] P. JEANQUARTIER, *Transformation de Mellin et développements asymptotiques*, L'Enseignement mathématique 25 (1979), 285–308.
- [21] M. LESCH AND N. PEYERIMHOFF, *An index theorem for manifolds with metric horns*, Comm. PDE **23** (1998), 649–684.
- [22] J.-M. LION AND J.-P. ROLIN, *Intégration des fonctions sous-analytiques et volumes des sous-ensembles sous-analytiques*, Preprint 1998.
- [23] M. NAGASE, *Hodge theory of singular algebraic curves*, Proc. Amer. Math. Soc. **108** (1990), 1095–1101.
- [24] R. M. RUIZ, *The basic theory of power series*, Vieweg 1993.
- [25] M. SPIVAK, *A comprehensive introduction to differential geometry, Vol. III-IV*, Publish or Perish, 1975.

AUTHOR'S ADDRESS:, INSTITUT FÜR MATHEMATIK, HUMBOLDT UNIVERSITÄT ZU BERLIN,
UNTER DEN LINDEN 6, 10099 BERLIN, GERMANY